

Instructions for diabetics before an endoscopy needing bowel preparation (**morning appointment**)

Endoscopy Unit
Information for Patients

Last reviewed: August 2026
Next review: August 2027
Leaflet number: 418 Version: 8

Introduction

If you are going to have a test called an endoscopy to look inside your bowel, you will need to stop eating (fast) for a while. This can make your blood sugar (glucose) levels drop

If you have diabetes and take medicine for it, you will need to change this according to the timing of your appointment to avoid having a 'hypo' (low blood sugar). This leaflet will help you to manage your diabetes to keep your blood sugar safe (between 4 and 11mmol/L).

Changing your medicine may upset your blood sugar levels. This should return to normal within 1 or 2 days after your procedure.

Please read the right instructions in this leaflet, depending on the type of medication you take:

- Follow the advice in **section 1** if you are only take **insulin**
- Follow the advice in **section 2** if you only take **tablets, or non-insulin injections**
- Follow the advice in **sections 1 and 2** if you take a **mix of insulin, tablets or any other injections**

If you have any questions about managing your diabetes for the test, please contact your usual diabetes care provider.

Please also read the instructions on taking the bowel preparation solution in your appointment pack.

Day before your procedure

- Have breakfast. Keep to the low residue diet that you are on. Please try and include white bread or white boiled rice with this meal.
- **Stop eating food after 1 pm the day before your procedure.** See the bowel preparation solution sheet for when to stop.
Drink lots of clear fluids like water, tea without milk. Follow the instructions on the accompanying bowel preparation solution sheet in your appointment pack.
- Check your blood sugar levels on a regular basis.
- You should have sugary drinks, if needed, to treat or avoid a 'hypo'.

Monitoring your blood sugar and treating a hypo

Hypoglycaemia (also known as a “hypo”) is when your blood sugar level falls too low. A blood sugar (glucose) below 4.0 mmol/L should be treated.

When you are on diabetes medication, a long period of fasting can make you more likely to get low blood sugar levels. Symptoms of hypoglycaemia include:

- blurred vision
- sweating a lot
- feeling anxious or angry
- tingling in mouth or fingers
- fast or heavy heartbeat
- odd behaviour (usually noticed by another person)
- trouble concentrating

It is important to check your blood sugar levels more often over this time (every 2 hours if you are worried).

What to do if you have a hypo whilst fasting:

If your blood sugar is under 4.0 mmol/L, you must break your fast. Have 2 to 5 glucose tablets.

Check your blood sugar level every hour, until your test. If during your hourly checks your blood sugar drops again below 4.0 mmol/L, you can take another glucose tablet.

If you have a hypo whilst preparing for your procedure, tell the staff when you arrive so we can monitor you more closely.

Section 1 - For patients taking insulin only

Name of insulin	Day before procedure	Day of procedure (AM appointment)	After procedure
Biphasic mixed insulins / 2 times daily			
Novomix 30 Humulinm3 Humalog mix 25 Humalog mix 50 Insuman comb 25 Insuman comb 50 Hypurin Porcine 30/70 mix	Reduce evening dose by a half.	a) Take half of your usual morning dose. b) Take your usual dose in the evening with food.	Take usual dose of insulin with food.
Basal/bolus insulin regime			
Bolus insulins: Actrapid Humulin S Insuman Rapid Novorapid Fiasp Humalog Apidra Lymjev Trurapi Hypurin Porcine Neutral	Do not take your evening dose of insulin.	a) Do not take your morning dose of insulin. b) Take usual dose at lunchtime with food. c) Take usual evening dose with food.	Take usual dose of insulin with food.
Basal insulins: Lantus Levemir Insulatard Humulinl Insuman basal Abasaglar Toujeo Tresiba Semglee Hypurin Porcine Isophane Xulutophy	If taken in the morning:		
	Take usual dose of insulin.	Take only half of your dose of insulin if taken insulin in the morning.	Take usual dose of insulin.
	If taken in the evening:		
	Reduce your insulin by a half.	Take usual dose of insulin.	

Section 2 - For patients taking tablets, or GLP-1 medicine or injections, for diabetes control

If you are taking tablets that are not mentioned here, please get medical advice from your usual diabetes care provider.

Medication	Day before procedure	Day of procedure	Day after procedure
Acarbose	Take as usual.	Do not take.	Take as usual.
Metformin	Take as usual.	Do not take.	Take as usual.
Repaglinide Nateglinide	Do not take evening dose.	Do not take morning or lunchtime dose.	Take as usual.
Gliclazide Glibenclamide Glipizide Glimepride Gliquidone	a) Take usual dose in the morning. b) Take half usual dose in the evening.	a) Do not take in the morning. b) Take as usual in the evening.	Take as usual.
Pioglitazone	Take as usual.	Take as usual.	Take as usual.
Dapagliflozin Canagliflozin Empagliflozin Ertugliflozin	Stop 4 days before procedure.	Do not take dose.	Take as usual.
Sitagliptin Saxagliptin Vildagliptin Alogliptin Linagliptin	Take as usual.	a) Do not take in the morning. b) Take as normal in the evening.	Take as usual.

GLP-1 injections and tablets			
Trulicity Exenatide (Byetta) Liraglutide Rybelsus Lixisenatide Semaglutide (Ozempic) Mounjaro	Take as usual	Take as usual	Take as usual

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Visit www.uhleicester.nhs.uk for maps and information about visiting Leicester's Hospitals.

اگر آپ کو یہ معلومات کسی اور زبان میں درکار ہیں، تو براہ کرم مندرجہ ذیل نمبر پر ٹیلی فون کریں۔
 ਜੇ ਤੁਸੀਂ ਇਹ ਜਾਣਕਾਰੀ ਕਿਸੇ ਹੋਰ ਭਾਸ਼ਾ ਵਿਚ ਚਾਹੁੰਦੇ ਹੋ, ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਹੇਠਾਂ ਦਿੱਤੇ ਗਏ ਨੰਬਰ 'ਤੇ ਟੈਲੀਫੋਨ ਕਰੋ।
 إذا كنت ترغب في الحصول على هذه المعلومات بلغةٍ أخرى، الرجاء الاتصال على رقم الهاتف الذي يظهر في الأسفل
 Aby uzyskać informacje w innym języku, proszę zadzwonić pod podany niżej numer telefonu
 જો તમને અન્ય ભાષામાં આ માહિતી જોઈતી હોય, તો નીચે આપેલ નંબર પર કૃપા કરી ટેલિફોન કરો.

If you would like this information in another language or format such as EasyRead or Braille, please telephone 0116 250 2959 or email uhl-tr.equalitymailbox@nhs.net

We are a research hospital and you may be asked to take part in research

Re-use of this leaflet is restricted by Creative Commons license

