

Diet and nutrition advice when you have gout

Dietetics and Nutrition service

Produced: May 2023

Review: May 2026

Information for Adult Patients

Leaflet number: 1400 Version: 1

What is gout?

Gout is a form of arthritis, affecting between 1 to 2 in every 100 adults in the UK. It is caused by having too much uric acid in your blood. This can lead to crystals of uric acid forming within or around the joints, causing swelling (inflammation). The symptoms include very bad pain, redness, swelling, discomfort and limited movement. Gout usually affects the big toe, but can also affect other areas.

How do uric acid crystals develop?

Purines are natural substances that are made by the body and also taken in with food. Purines are broken down into uric acid when food is digested. Uric acid is normally removed in the urine by the kidneys.

If not enough purine is removed, or if you have a high intake of purine rich foods, the levels of uric acid can build up in the blood. These form solid crystals around the joints. If left untreated, high levels of uric acid can increase your risk of heart disease.

Risk factors

You may be more likely to get gout if you have:

- A family history of gout.
- Obesity, high blood pressure or diabetes.
- Kidney disease.
- A high alcohol intake (often drinking more than 14 units a week).

**Health information and support is available at www.nhs.uk
or call 111 for non-emergency medical advice**

Visit www.leicestershospitals.nhs.uk for maps and information about visiting Leicester's Hospitals
To give feedback about this information sheet, contact InformationForPatients@uhl-tr.nhs.uk

How can gout be managed?

Your symptoms can be managed with medical treatments and lifestyle changes. This leaflet focuses on changes to your diet. If you would like to know more about medications for gout, please talk to your doctor, or visit www.ukgoutsociety.org.

Healthy weight

Having a healthy lifestyle can help to reduce gout flares and symptoms. Being overweight or obese may increase your risk of gout or make the symptoms worse. If you would like help to lose weight, talk to your GP or Dietitian, or visit www.nhs.uk/better-health/lose-weight/.

Alcohol

Drinking a lot of alcohol can make your symptoms worse, or more likely to happen more often.

Current national guidelines advise that both men and women should avoid drinking more than 14 units of alcohol in a week on a regular basis. (For example, a typical 'single' measure of vodka (25 ml) is 1 unit, a pint of 4% lager is 2.3 units).

All types of alcohol can increase your risk of repeated gout attacks.

For more information on reducing the amount of alcohol you drink, please talk to your GP or visit: www.drinkaware.co.uk.

Diet

Research around gout and diet is showing mixed results and at the moment there is not enough evidence to suggest that any specific type of diet stops flare ups. Current clinical guidelines recommend that you aim to have a healthy, balanced diet.

Some studies suggest that a low purine diet and eating less calories may help manage gout. Please talk to your Dietitian if you are not sure which approach is right for you.

Low-purine diet


Purines are broken down into uric acid. They are part of a normal diet, and cannot be avoided completely. You can try to reduce the amount of purines in your diet by eating less of the 'high purine' foods (see table below).

Tip: focus on the portion sizes and frequency, you do not have to completely remove any of these foods from your diet.

Purine content	Food	Portion size
High Avoid eating regularly	Offal - liver and kidneys, heart and sweetbreads	85-90g
	Game - pheasant, rabbit, venison	90g
	Oily fish - anchovies, herring, mackerel, sardines, sprats, whitebait, trout	140g, palm of hand
	Seafood - especially mussels, crab, shrimps and other shellfish, fish roe, caviar	85g
	Meat and yeast extracts - Marmite, Bovril, gravy granules	Marmite: 8g Bovril: 12 g Gravy: 50ml
Medium Eat smaller portions, less frequently (in moderation)	Meat - beef, lamb, chicken, pork	90g, palm of hand
	Poultry - chicken and duck	90g, palm of hand
	Dried peas, beans and legumes - baked beans, kidney beans, soya beans and peas	150g or 4 tablespoons
	Mushrooms and mycoprotein (for example Quorn™)	100g or 4 tablespoons
	Some vegetables - asparagus, cauliflower, spinach	80g
	Wholegrains - bran, oat bran, wholemeal bread	One medium slice of bread (34-36g) 30-40g of cereal (3 tablespoons)
Low No restrictions	Dairy - preferably low or reduced fat	A small matchbox-sized piece of cheese 200ml glass of milk 125 to 150g yoghurt
	Eggs	120g – 2 medium
	Bread and cereals (except wholegrain)	1 medium slice of bread (34 to 36g) 30 to 40g of cereal (3 tablespoons)
	Pasta and noodles	75g uncooked/150g cooked
	Fruit Vegetables (see moderate purine list)	80g, for example 1 medium apple, banana or pear, ½ a pepper or 7 cherry tomatoes

Healthy eating

Making sure that you eat a healthy, balanced diet is key to prevent a lack of essential vitamins and minerals (micronutrient deficiencies) that could make you unwell. Use the Eatwell guide to help you make good dietary choices and maintain a healthy body weight. You can also use the layout of the plate in the picture to help you balance your meals.



Source: Public Health England in association with the Welsh Government, Food Standards Scotland and the Food Standards Agency in Northern Ireland

© Crown copyright 2016

- Aim to have 5 portions of fruit and vegetables per day (80g is 1 portion).
- Try to limit how much salt (sodium) you eat (no more than 6g a day, equal to 1 teaspoon).
- Try to reduce how much sugar you eat by cutting back on added sugars, for example from sweetened drinks, sweets and chocolate.
- Use unsaturated fats (for example olive oil) and try to reduce saturated fats (for example from fatty cuts of meat, lard).
- You could reduce the amount of fat you eat by swapping full fat dairy for reduced or low fat options.
- Have lean proteins such as fish, beans, lentils, nuts, seeds and pulses.

Fibre

Some studies suggest that having a high fibre diet (30 grams of fibre a day) can help to reduce swelling caused by gout.

Which foods are high in fibre?

Foods	Portion size	Grams of fibre
Ready Brek	30g	2g
Porridge*	30g	4g
Wholemeal bread*	Thick slice	3g
Weetabix*	1 biscuit	2g
Wholemeal pasta*	100g	4g
Apple	Medium	2g
Pear	Medium	5g
Strawberries	100g	4g
Baked beans*	200g	7g
Broccoli	100g	3g
Raw peppers	80g	2g
Brussel sprouts	3 sprouts	3g
Peanuts	30g	3g
Jacket potato with skin on	150g	3g
Wholegrain rice*	100g	2g
Quorn™ pieces*	50g	4g

*Eat in moderation if you are on a low purine diet.

Vitamin C

There is some limited evidence to suggest that eating foods that have more Vitamin C, for example from cherries, sour cherry juice or a supplement can reduce the risk of gout flare-ups.

If you are thinking of taking an over the counter supplement, get advice from your doctor or pharmacist.

Fluid

Drinking more fluid can help to remove uric acid through the kidneys. This is very important when you are having a flare-up. You should aim to drink at least 8 large glasses of fluid a day. If you have kidney disease or are taking water tablets (diuretics, for example furosemide) you should talk to a doctor before you increase your fluid intake.

If you need any more information please talk to your Dietitian.

Dietetics and Nutrition Service

University Hospitals of Leicester NHS Trust

Dietitian: _____

Telephone number: 0116 258 5400

اگر آپ کو یہ معلومات کسی اور زبان میں درکار ہیں، تو براہ کرم مندرجہ ذیل نمبر پر ٹیلی فون کریں۔
على هذه المعلومات بلغةٍ أخرى، الرجاء الاتصال على رقم الهاتف الذي يظهر في الأسفل
જો તમને અન્ય ભાષામાં આ માહિતી જોઈતી હોય, તો નીચે આપેલ નંબર પર કૃપા કરી ટેલિફોન કરો
ਜੇ ਤੁਸੀਂ ਇਹ ਜਾਣਕਾਰੀ ਕਿਸੇ ਹੋਰ ਭਾਸ਼ਾ ਵਿਚ ਚਾਹੁੰਦੇ ਹੋ, ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਹੇਠਾਂ ਦਿੱਤੇ ਗਏ ਨੰਬਰ 'ਤੇ ਟੈਲੀਫੋਨ ਕਰੋ।
Aby uzyskać informacje w innym języku, proszę zadzwonić pod podany niżej numer telefonu

Previous reference:

If you would like this information in another language or format such as EasyRead or Braille, please telephone 0116 250 2959 or email equality@uhl-tr.nhs.uk