

# Therapy after tendon injury to your finger - SAM exercises

## Occupational Therapy

Information for Patients

Produced: September 2023

Review: September 2026

Leaflet number: 1434 Version: 1

### Introduction

- Tendons are cords of tissue that attach muscles to the bones, these structures all work together to make the joints in your hand move.
- There are 2 types of tendon in your hand. Extensor tendons run along the top of the hand. They help to straighten fingers. Flexor tendons run across the palm of the hand. They help the fingers to bend.
- You have had an injury to your finger.
- Damage to a tendon can occur through any type of trauma causing the tendon to be cut or crushed.
- It takes at least 12 weeks after the tendon is repaired for it to regain full strength.
- Tendons need a long period of therapy because the repair will be very weak at first. Too much stress, too early, through the repair will cause it to break and fail.
- You have cut the tendon(s) in your hand and they have been repaired by the surgeon.
- Tendon repair involves making a cut in your hand, finding the ends of the tendon, and then stitching them back together. .


### What is SAM?

After your surgery you will wear a splint and do a number of exercises every day. The exercises are called short arc motion or SAM for short.

**Health information and support is available at [www.nhs.uk](http://www.nhs.uk)  
or call 111 for non-emergency medical advice**

Visit [www.leicestershospitals.nhs.uk](http://www.leicestershospitals.nhs.uk) for maps and information about visiting Leicester's Hospitals  
To give feedback about this information sheet, contact [InformationForPatients@uhl-tr.nhs.uk](mailto:InformationForPatients@uhl-tr.nhs.uk)

## Your splints



A splint will be made for you to keep your finger straight. This protects your tendon repair by stopping you from bending your fingers too far.

It is important you **do not remove** your splint.

If you bend your fingers more than allowed, your tendon repair is at risk of breaking. If this occurs you will need to have another operation to repair it.

For 4 weeks you will wear the splint at all times. Please do not use your hand for everyday activities during this time.

## Exercises


Please do these exercises every 2 hours

### Exercise 1



- Bend the main knuckles of the hand as far as you can.
- Return to fully straight.
- Repeat 15 times

## Exercise 2



- Remove the end strap.
- Support the middle joint with the blocking splint.
- Bend the tip of your finger as far as you can.
- Repeat 15 times.

## Exercise 3

**For this exercise please keep your wrist bent gently forwards**



- Remove the top 2 straps.
- Place the exercise splint you have been given under your finger.
- With your wrist slightly bent forwards bend the finger to the exercise splint and then back to fully straight.
- Repeat 15 times

## General advice

### Activity:

- Do not use your hand for **any** activity for 4 weeks.
- Only do the exercises you have been advised to do by your therapist. It does not help to do more exercises than advised.
- At 4 weeks you will be able to do light tasks. Your therapist will tell you which activities are safe to do.
- You will be told not to carry out any heavy lifting or forceful gripping until at least 12 weeks after your operation.
- Specific advice will be given if you are in employment.
- You are advised not to drive for at least 8 weeks. Your therapist will give you further guidance.

### Swelling:

- You may have some swelling in your hand after your operation. This is quite normal.
- To reduce this it is important to raise (elevate) your hand above your heart level. This can be done whilst sitting or lying by resting your hand across your chest/shoulder or by using pillows.
- Do not walk around with your hand hanging down at your side for long periods during the first 4 weeks after your operation.

## Complications

If your wound becomes hot, red, very painful or leaks fluid it could be infected, please call your GP as soon as possible.

If you notice that you cannot move your finger(s) in the way you did before or if you have felt a 'snapping', your tendon repair could have broken. If this occurs you should contact the surgical team at the hospital as soon as possible. If you are unable to contact the surgical team, contact your Therapist.

Your tendons may become stuck to surrounding tissues. This may mean you have less range of movement in the finger(s). Your therapist will advise you about this.

Damage can occur to the nerves in your finger (s) either at the time of your injury or during the surgery to repair your tendon. If this occurs it will affect the feeling in your finger(s). This may get better gradually or you may always have less feeling. Your therapists will talk to you about this.

## Contact details

Name of Occupational Therapist.....

Telephone Number.....

اگر آپ کو یہ معلومات کسی اور زبان میں درکار ہیں، تو براہ کرم مندرجہ ذیل نمبر پر ٹیلی فون کریں۔  
على هذه المعلومات بلغة أخرى، الرجاء الاتصال على رقم الهاتف الذي يظهر في الأسفل  
જો તમને અન્ય ભાષામાં આ માહિતી જોઈતી હોય, તો નીચે આપેલ નંબર પર કૃપા કરી ટેલિફોન કરો  
ਜੇ ਤੁਸੀਂ ਇਹ ਜਾਣਕਾਰੀ ਕਿਸੇ ਹੋਰ ਭਾਸ਼ਾ ਵਿਚ ਚਾਹੁੰਦੇ ਹੋ, ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਹੇਠਾਂ ਦਿੱਤੇ ਗਏ ਨੰਬਰ 'ਤੇ ਟੈਲੀਫੋਨ ਕਰੋ।  
Aby uzyskać informacje w innym języku, proszę zadzwonić pod podany niżej numer telefonu

Previous reference:

If you would like this information in another language or format such as EasyRead or Braille, please telephone 0116 250 2959 or email [equality@uhl-tr.nhs.uk](mailto:equality@uhl-tr.nhs.uk)