

# Hip fracture

Emergency Department

Last reviewed: June 2023

Information for Patients

Next review: June 2026

Leaflet number: 373 Version: 2

## What is a hip fracture?

The hip is a ball and socket joint. The socket is in your pelvis and the ball is the head of your thigh bone (femur).

When you break or crack a bone this is called a fracture. A break in the top of your thigh bone, near to where the ball and socket meet, is called a hip fracture.

## Why do people get hip fractures?

The most common reason why people get hip fractures is due to a fall. Older people are at higher risk of falling over as we tend to lose our strength and balance as we grow older.

As we get older, our bones can become thinner, which is a condition called osteoporosis. This means that our bones are weaker and more likely to break if we injure ourselves.

## How do we treat a hip fracture?

Hip fractures are usually treated with an operation. The main aims of the operation are to reduce the pain from the hip fracture and to enable you to get back on your feet as soon as possible.

The type of operation you need depends on which part of your hip has been broken. Your doctor will talk to you about which operation is best for you.

Types of operation:

- Internal repair - The surgeon uses pins, plates, screws or rods to hold the broken bones in place to allow them to heal.

**Health information and support is available at [www.nhs.uk](http://www.nhs.uk)  
or call 111 for non-emergency medical advice**

Visit [www.leicestershospitals.nhs.uk](http://www.leicestershospitals.nhs.uk) for maps and information about visiting Leicester's Hospitals  
To give feedback about this information sheet, contact [InformationForPatients@uhl-tr.nhs.uk](mailto:InformationForPatients@uhl-tr.nhs.uk)

- Partial hip replacement – The head of your thigh bone is replaced with metal implants.
- Total hip replacement - The head of your thigh bone and the socket in your pelvis are replaced with metal implants

## How can we help reduce the pain?

In the Emergency Department, you will be offered pain killers. We often use a few different pain killers together, as this helps most with pain relief. You may be offered an injection of painkilling medication (local anaesthetic) into the groin, called a nerve block. This will help to reduce the pain for quite a few hours so that you are more comfortable moving to the ward.

It is important to take pain medication regularly as they take time to work and you may not feel the relief straight away.

## When will I have my operation?

We aim to do the operation within 36 hours of you being admitted to hospital with a broken hip. Sometimes this is not possible, for example if you have a medical condition that needs to be treated before you have surgery.

## What happens after the operation?

You will work with a team of physiotherapists and occupational therapists. This will allow you to start putting weight through your hip the day after the operation. Our aim is to get you walking early so that you can regain your independence sooner. This will also help avoid complications such as blood clots in the lungs and legs. You may also be offered injections of blood thinners to help stop clots from developing.

Walking can be very sore to start with, which is normal. It is important that you are involved in your recovery and we will encourage you to do this. The occupational therapist may suggest changes to your home and day to day activities to help you regain your independence.

## When can I go home?

Our average length of stay for patients who have had a hip fracture is 11 days. This may be longer - particularly in patients who have lots of medical problems. The team looking after you will keep you updated with your progress. They will work with you and a relative or carer, with your permission, to get you home when it is safe to do so.

## How can I prevent another fracture?

You should go to your GP (general practitioner) to see if there is a reason for your fall or fracture. If you have thinning of the bones (osteoporosis), you may need treatment with vitamin D, calcium supplements, or tablets that can help strengthen your bones. It is important to keep active, have a regular medication review with your GP and annual eye checks. You should eat a healthy balanced diet rich in calcium, vitamin D and protein. It is important to reduce or stop smoking. A fracture can take longer to heal if you smoke. We can support you to stop and also prescribe nicotine replacement.

## How long will it take to recover?

It can take quite a few months to recover from a hip fracture. Every patient and injury is different and so is the healing time. Follow the advice of your team for a speedy recovery.

اگر آپ کو یہ معلومات کسی اور زبان میں درکار ہیں، تو براہ کرم مندرجہ ذیل نمبر پر ٹیلی فون کریں۔  
على هذه المعلومات بلغةٍ أخرى، الرجاء الاتصال على رقم الهاتف الذي يظهر في الأسفل  
જો તમને અસ્ય ભાષામાં આ માહિતી જોઈતી હોય, તો નીચે આપેલ નંબર પર કૃપા કરી ટેલિફોન કરો  
ਜੇ ਤੁਸੀਂ ਇਹ ਜਾਣਕਾਰੀ ਕਿਸੇ ਹੋਰ ਭਾਸ਼ਾ ਵਿਚ ਚਾਹੁੰਦੇ ਹੋ, ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਹੇਠਾਂ ਦਿੱਤੇ ਗਏ ਨੰਬਰ 'ਤੇ ਟੈਲੀਫੋਨ ਕਰੋ।  
Aby uzyskać informacje w innym języku, proszę zadzwonić pod podany niżej numer telefonu

Previous reference:

If you would like this information in another language or format such as EasyRead or Braille, please telephone 0116 250 2959 or email [equality@uhl-tr.nhs.uk](mailto:equality@uhl-tr.nhs.uk)