

Aftercare following trabeculectomy surgery for glaucoma

Department of Ophthalmology

Information for Patients

Last reviewed: January 2026

Next review: January 2029

Leaflet number: 1141 Version: 3

What will happen after your surgery?

- After your operation we will put a clear plastic shield over your eye. This is to protect it from accidental knocks.
- You will be able to go home after the operation.
- We will give you 2 sets of eye drops. The eye drops are an antibiotic (like chloramphenicol) and an anti-inflammatory steroid (like dexamethasone or predforte).
- **Steroid eye drop:** You will need to use the steroid eye drop every 2 hours during the day. The steroid eye drops need to be used for a few months in order to help stop scarring. The surgical team will tell you if you need to use this eye drop for longer. The dose will be changed by your doctor at your clinic visits, based upon how your eye is healing.
- **Antibiotic eye drop:** You will need to use this 4 times a day. This is often for 2 weeks only. You must keep using any eye drops that you were using in your other eye.
- You will need to come back the next day so your consultant can check your eye after the operation. At this visit, we will remove the eye shield for you. We will tell you about any other appointments you may need after your operation. We will check your eye's recovery.

Health information and support is available at www.nhs.uk
or call 111 for non-emergency medical advice

Visit www.uhleicester.nhs.uk for maps and information about visiting Leicester's Hospitals
To give feedback about this information sheet, contact uhl-tr.informationforpatientsmailbox@nhs.net

How will my eye feel afterwards?

- You may get blurred or double vision. This can last for up to 1 month.
- Your eye may feel uncomfortable and gritty. You can take painkillers you take for a headache like paracetamol to help with this.
- The eye can look red and swollen. We will give you anti-inflammatory eye drops to help with this. Your discharge nurse will tell you how often to use these.
- In some cases we may need to give you an injection to help stop scar tissue forming. We can give this injection in the outpatient clinic.
- Your eye should slowly get better over the weeks after the operation. However, if you notice any of the following:
 - increased redness
 - increased swelling
 - sudden complete loss of sight
 - a drop in your vision
 - pain in the eye
 - yellow sticky discharge

You should contact the Ophthalmic Suite or Eye Casualty using the numbers on the last page of this leaflet.

After surgery you should:

- Wear the eye shield until the morning after your operation.
- Wear the eye shield in bed for the next 14 nights. Use surgical tape to hold the eye shield in place. You can buy this from pharmacies and supermarkets.
- Wash the eye shield daily with hot soap and water.
- If you wish, you can wear sunglasses or normal glasses during the day.
- If your eye is sticky in the mornings, you can bathe it with cooled boiled water and cotton wool. If you are concerned about this then please contact us.
- Cooking, watching TV and reading will not cause any harm.
- Avoid energetic activity like contact sports and swimming for 1 month. Avoid heavy gardening and any lifting for 4 weeks after your operation.

After surgery do not:

- Do not drive until you have checked with a doctor, nurse or your optician that it is safe to do so.
- Do not rub or touch your eye.
- Do not get soap and water in your eye when washing your face and hair.

Using your eye drops


Your old glaucoma drops in the operated eye should be stopped and the following drops should be used on your **operated eye only**. You should continue any eye drops you are using to the non operate eye as before.

Start this the morning after surgery:

Eye drop	Time	Action
Chloramphenicol eye drops (to keep in fridge)	<ul style="list-style-type: none"> • 1 drop, • 4 times a day, • for 2 weeks, then stop. 	Antibiotic
Dexamethasone eye drops	<ul style="list-style-type: none"> • 1 drop, • every 2 hours during the day, • for 2 weeks, reducing after this as per the advice in clinic. The usual total course of these drops is three months. 	Anti-inflammatory

How to apply your eye drops

- Always wash your hands before you start.
- Sit or lie with your head tilted backwards.
- Gently pull the lower lid down with one finger.
- Holding the bottle on the bridge of your nose or on your forehead, squeeze one drop into the eye.
- Close your eye and gently blot excess drops with a clean tissue.
- If you think you may have missed your eye with the drops it is safe to put another drop in.
- When putting the eye drops into your eye, do not let the bottle tip to touch the eye.
- Do not let other people use your eye drops.
- Throw away the drops when you have finished the course of treatment.



Eye drop chart for chloramphenicol eye drops

Please use the chart below to record when you have applied your chloramphenicol eye drops to your operated eye.

	Breakfast	Lunch	Dinner	Bedtime
Day 1				
Day 2				
Day 3				
Day 4				
Day 5				
Day 6				
Day 7				
Day 8				
Day 9				
Day 10				
Day 11				
Day 12				
Day 13				
Day 14				

Eye drop chart for Dexamethasone eye drops

Please use the chart below to record when you have applied your Pred Forte eye drops to your operated eye.

	8:00am	10:00am	12:00pm	2:00pm	4:00pm	6:00pm	8:00pm
Day 1							
Day 2							
Day 3							
Day 4							
Day 5							
Day 6							
Day 7							
Day 8							
Day 9							
Day 10							
Day 11							
Day 12							
Day 13							
Day 14							



Contact details

If you have any questions or concerns, please contact:

The Ophthalmology Suite:

0116 258 6248 or 0116 258 6850

Monday to Friday from 8:30am to 7:30pm

Eye Casualty department:

0116 258 6273

Monday to Friday from 8:00am to 6:00pm

Weekend and Bank Holidays 8:00am to 12:30pm

At other times please contact your **nearest Emergency Department** for urgent concerns.

اگر آپ کو یہ معلومات کسی اور زبان میں درکار ہیں، تو براہ کرم مندرجہ ذیل نمبر پر ٹیلی فون کریں۔
على هذه المعلومات بلغةٍ أخرى، الرجاء الاتصال على رقم الهاتف الذي يظهر في الأسفل
જો તમને અન્ય ભાષામાં આ માહિતી જોઈતી હોય, તો નીચે આપેલ નંબર પર કૃપા કરી ટેલિફોન કરો

ਜੇ ਤੁਸੀਂ ਇਹ ਜਾਣਕਾਰੀ ਕਿਸੇ ਹੋਰ ਭਾਸ਼ਾ ਵਿਚ ਚਾਹੁੰਦੇ ਹੋ, ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਹੇਠਾਂ ਦਿੱਤੇ ਗਏ ਨੰਬਰ 'ਤੇ ਟੈਲੀਫੋਨ ਕਰੋ।
Aby uzyskać informacje w innym języku, proszę zadzwonić pod podany niżej numer telefonu

If you would like this information in another language or format such as EasyRead or Braille, please telephone 0116 250 2959 or email uhl-tr.equalitymailbox@nhs.net