

Anaphylaxis action plan - for use with Emerade® adrenaline auto-injector pen

Allergy Service

Produced: March 2022

Review: March 2025

Information for Patients

Leaflet number: 318 Version: 2

Name:	Date of birth:
NHS number:	
Emergency contact:	
Consultant:	
Plan prepared by:	
Date:	

Follow the action plan over the page if you have any **ONE** of these symptoms of a **severe allergic reaction (anaphylaxis)**:

- Airway difficulties:** Swelling of tongue
Swelling / tightness in throat
Difficulty talking / hoarse voice
- Breathing:** Difficult / noisy breathing
Wheeze or persistent cough
- Circulation:** Persistent dizziness or collapse
Loss of consciousness

Health information and support is available at www.nhs.uk
or call 111 for non-emergency medical advice

Visit www.leicestershospitals.nhs.uk for maps and information about visiting Leicester's Hospitals
To give feedback about this information sheet, contact InformationForPatients@uhl-tr.nhs.uk

Action plan for a severe reaction

1. **Lie flat** with your feet up until help arrives. If breathing is difficult you can sit, but do not stand or walk.
2. **Use your Emerade[®]**.
3. **Call 999** for an ambulance and say you are having anaphylaxis (“ana-fil-ax-is”).
4. If there is no improvement after 5-15 minutes use your second Emerade[®] in the other leg.
5. Take antihistamines as prescribed (if not already taken). If you are asthmatic use reliever inhaler.

If in doubt or rapidly deteriorating, use your Emerade[®].

Medical observation in hospital is recommended after anaphylaxis.

Always keep your 2 Emerade pens with you at all times.


Follow the action plan below if you have symptoms of a mild or moderate allergic reaction, such as:

- Swelling of lips, face, or eyes.
- Hives or welts.
- Tingling mouth.
- Abdominal pain/ vomiting.

Action plan:

1. Locate your Emerade[®].
2. Take antihistamines as prescribed.
3. Phone family/ emergency contact.
4. **Watch for worsening symptoms; if in doubt or rapidly deteriorating use your Emerade[®].**

How to give Emerade[®]



REMOVE NEEDLE SHIELD



PRESS AGAINST THE OUTER THIGH



HOLD FOR 5 SECONDS

Massage the injection site gently

Store Emerade[®] in a cool dark place at room temperature; keep out of direct sunlight and avoid extreme temperatures (e.g. fridge). For more information and to register for a free expiry alert service visit:

www.emerade-bausch.co.uk

For more information on anaphylaxis visit:

- www.allergyuk.org



- www.anaphylaxis.org.uk



If you would like this information in another language or format such as EasyRead or Braille, please telephone 0116 250 2959 or email equality@uhl-tr.nhs.uk